



## lecanemab-irmb

LEQEMBI

100 mg/mL

## Dose Preparation


**NATIONAL  
INFUSION CENTER  
ASSOCIATION**

## Order Details

## Patient Weight:

(enter patient weight in kg  
and click on Tab key)

 kg

## Dose @ 10 mg/kg

 mg

## Dose Volume:

 mL

## Diluent Volume:

0.9% Sodium Chloride

 mL

## Total Volume:

(Dose Volume + Diluent  
Volume)

 mL

## Dose Range

500 mg / 5 mL

vials needed

200 mg / 2 mL

vials needed

500 mg

1

0

501 - 700 mg

1

1

701 - 900 mg

1

2

901 - 1000 mg

2

0

1001 - 1200 mg

2

1

1201 - 1400 mg

2

2

1401 - 1500 mg

3

0

## Dose Preparation Instructions

- Calculate the number of LEQEMBI (lecanemab-irmb) vials needed for the dose (see above). Each vial contains a concentration of 100 mg / mL.
- Remove LEQEMBI (lecanemab-irmb) vial(s) from refrigerated storage and allow to warm to room temperature. Do not expose to direct heat.
- Visually inspect the LEQEMBI (lecanemab-irmb) diluted solution for particles or discoloration prior to administration. Do not use if it is discolored, or opaque or foreign particles are seen.
- Remove the flip-off cap from the vial. **Disinfect and allow to air dry.** Using aseptic technique, insert the sterile syringe needle into the vial through the center of the rubber stopper.
- Slowly withdraw  mL of LEQEMBI (lecanemab-irmb) from the vial. Each vial is for single-use only; discard any unused portion.
- Disinfect injection port of infusion bag containing 250 mL of 0.9% Sodium Chloride Injection and allow to air dry. Using aseptic technique, slowly inject LEQEMBI (lecanemab-irmb) into infusion bag. Avoid creating turbulence or foaming.
- Gently invert the infusion bag containing the LEQEMBI (lecanemab-irmb) diluted solution to mix completely. **Do not shake. Do not freeze.**
- After dilution, immediate use is recommended.** If not administered immediately, store LEQEMBI (lecanemab-irmb) refrigerated for up to 4 hours, or at room temperature for up to 4 hours.

lecanemab-irmb

LEQEMBI

Dose Administration


**NATIONAL  
INFUSION CENTER  
ASSOCIATION**

## Order Details

Dose:  mgDose Volume:  mLDiluent Volume:  
0.9% Sodium Chloride  mLTotal Volume:  
(Dose Volume + Diluent  
Volume)  mL

## Order Details

Administration Time 1 hour

Filter 0.2 micron  
in-line filterFlush Yes  
*Flush infusion line to insure all  
drug is administered.*Infusion Rate  mL/hr

## Administer by Intravenous Infusion

Attach the infusion set (containing a terminal low-protein binding 0.2 micron in-line filter) to the IV bag. Prime the infusion set.

Infuse the entire volume of the LEQEMBI (lecanemab-irmb) diluted solution intravenously over approximately one hour.

Flush infusion line to ensure all LEQEMBI (lecanemab-irmb) is administered.

**Monitor for signs and symptoms of an infusion reaction.**

The infusion rate may be reduced, or the infusion may be discontinued, and appropriate therapy administered as clinically indicated.

Consider pre-medication at subsequent dosing with antihistamines, non-steroidal anti-inflammatory drugs, or corticosteroids, if signs or symptoms of an infusion-related reaction are observed.